


# Environmental Weed Risk Assessment results for

## Cucurbita pepo L.

Assessors Name/s RR




Maps: Current Australian distribution [above left], predicted preferred climate [above right] and global distribution [bottom].

Common Name/s: marrow, pumpkin

|                           |             |                      |             |
|---------------------------|-------------|----------------------|-------------|
| <b>Invasiveness Score</b> | <b>2.96</b> | <b>Impacts Score</b> | <b>0.50</b> |
|---------------------------|-------------|----------------------|-------------|

|           | Climate Score | Final FFI Score | Environmental Weed Risk Rating |
|-----------|---------------|-----------------|--------------------------------|
| Kimberley | 7             | 10.37           | Negligible                     |
| Pilbara   | 2             | 2.96            | Negligible                     |
| Gascoyne  | 2             | 2.96            | Negligible                     |
| Southwest | 8             | 11.85           | Negligible                     |



| FINAL FFI Score | Weed Risk  | Interpretation - Invasive Species Weed Assessor               |
|-----------------|------------|---|
| > 235           | Very High  | Use not supported, significant environmental risks            |
| 112 - 235       | High       | Use not recommended, any spread to be reported and controlled |
| 50 - 111        | Medium     | Can be used, monitor, manage and report any offsite movement  |
| 18 - 49         | Low        | Can be used with minimal concern                              |
| < 18            | Negligible | Can be used without concern                                   |

For more details on the CRC Future Farm Industries system see:

Stone, L., Munday, C. and Bettink, K. (2012). Environmental Weed Risk Assessment Protocol. Future Farm Industries CRC (20 pp.). <http://www.futurefarmonline.com.au/about/weedrisk.htm>