
Environmental Weed Risk Assessment results for

Leucaena collinsii Britton & Rose

Synonym/s:

Assessors Name/s: Rod Randall

Assessment Date: Date Last Modified: 8/06/2018



Maps: Current Australian distribution [above left], predicted preferred climate [above right] and global distribution [bottom].

Common Name/s: leadtree

Invasiveness Score

5.56

Impacts Score

6.50

Final FFI Score = Invasiveness Score x Impacts Score x Climate Score

| | Climate Score | Final FFI Score | FFI Environmental Weed Risk Rating | Region Boundaries |
|-----------|------------------|--------------------|---------------------------------------|--|
| Kimberley | 1 | 36.11 | Negligible - Low | FOUR BASIC CLAMATIC RECIONS ASSESSED MINISTRUT ANALYA MACINE MACIN |
| Pilbara | 0.5 | 18.06 | Negligible - Low | MALE OFFICE AMERICAN |
| Gascoyne | 0.5 | 18.06 | Negligible - Low | CARACTORY WINDOWS MARKET SAMPAUL MAR |
| Southwest | 0.5 | 18.06 | Negligible - Low | OCCUPATION TOLLOWS BOALDER |

| FINAL FFI Score | Weed Risk | Interpretation - Invasive Species Weed Assessor |
|-----------------|------------------|--|
| > 235 | Very High | Use not recommended, requires a management plan, significant risks |
| 112 - 235 | High | Use not normally recommended, requires a risk management plan |
| 50 - 111 | Medium | Can be used, monitor, manage and report any offsite movement |
| 0 - 49 | Negligible - Low | Can be used with minimal concern |
| | | |

For more details on the CRC Future Farm Industries system see:

Stone, L., Munday, C. and Bettink, K. (2012). Environmental Weed Risk Assessment Protocol. Future Farm Industries CRC (20 pp.). http://www.futurefarmonline.com.au/about/weedrisk.htm